

**ADDENDUM TO: TORSION AND DEFORMATION
OF CONTACT METRIC STRUCTURES
ON 3-MANIFOLDS**

(Tôhoku Math. J. 39 (1987), 365–372)

SAMUEL I. GOLDBERG¹ AND GABOR TOTH

(Received December 17, 1987)

Chern and Hamilton [1] introduced the torsion invariant $|\tau|$ on a compact contact Riemannian 3-manifold (M, g) , where $\tau = L_{X_0}g$, and they conjectured that for fixed contact form ω , with X_0 inducing a Seifert foliation, there exists a complex structure $\varphi|_B$ on $B = \ker \omega$ with the property that the Dirichlet energy

$$(1) \quad E(g) = \frac{1}{2} \int_M |\tau|^2 \operatorname{vol}(M, g)$$

is critical over all CR structures. A CR structure on a manifold is a contact structure together with a complex structure on B . Since $\dim M = 3$, B has dimension 2, so a complex structure on B is equivalent to a conformal structure, i.e., knowing how to rotate through 90° . Hence, a Riemannian metric on a contact 3-manifold gives rise to a CR structure.

Let M be a contact manifold with a fixed contact form ω . Denote the space of all associated Riemannian metrics to the contact form ω by $\mathcal{M}(\omega)$. Since $E(g)$ vanishes, if and only if τ vanishes, this implies that X_0 is a Killing vector field. Can E have a critical point which is not a zero of E ? Let g be a point of \mathcal{M} , and $\{g(t)\}$ be a curve in \mathcal{M} with $g(0) = g$. Tanno [5] showed that g is a critical point of E , i.e., $(dE/dt)(0) = 0$, if and only if

$$(2) \quad \nabla_{X_0}\tau = 2\tau \cdot \varphi.$$

Thus, following [1], $E(g)$ is critical over all CR structures if and only if (2) is satisfied. This is different from the condition $\nabla_{X_0}\tau = 0$ incorrectly obtained in [1, Theorem 5.4]. The reason for this discrepancy is given by Tanno in [5, p. 15]. The statement of the Theorem in [3] should therefore be modified by replacing the phrase “and critical torsion” by the phrase “such that $\nabla_{X_0}\tau = 0$ ”. (The condition $\nabla_{X_0}\tau = 0$ is equivalent to the statement that the sectional curvature of all planes at a given point perpendicular to B are equal (see [1] and [5]).)

In the sequel, a critical point of E will be called a *critical metric*. If g is a critical metric, then by [3, Proposition 1, formula (ii)], $\nabla_{X_0}\tau = -2\psi$, where $\psi(X, Y) =$

¹ Supported by the Natural Sciences and Engineering Research Council of Canada.

$g((L_{X_0}\varphi)X, Y)$. Formula (4) of [3] with $n=1$ and $b=a^2-a$ then reduces to

$$(3) \quad \tilde{S} = S + 2(1-a)g + 2(a-1)(a+2)\omega \otimes \omega$$

which is somewhat surprising since the torsion terms cancel. (This formula is the same as the one obtained in [2, p. 654], where it is assumed that X_0 is a Killing vector field.) With several important modifications, the method of proof of the Theorem in [3] also gives rise to the following:

THEOREM. *Let M be a compact orientable 3-manifold with contact metric structure $(\varphi, X_0, \omega, g)$ where g is critical. If there exists a constant a , $0 < a < 1$, such that $c < 2a$, and if*

$$(4) \quad |\sigma|^2 < 2 \left(a^2 - \frac{c^2}{4} \right) \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a - c \right),$$

then M admits a contact metric of positive Ricci curvature. If, in addition, M is simply connected, it is diffeomorphic with the standard 3-sphere S^3 .

PROOF. We show that $\tilde{S} > 0$ at each $x \in M$. To this end, we determine the matrix \tilde{S} in (3) with respect to a suitable adapted basis $\{E, \varphi E, X_0\}$ of $T_x M$, and compute the respective subdeterminants along the main diagonal. Assume $\sigma \neq 0$. (If $\sigma = 0$, the same argument applies.) Since σ is a linear form on B , $\ker \sigma \neq 0$. Choose E such that $\sigma(E) = 0$ and $\sigma(\varphi E) = |\sigma|$. Then,

$$\tilde{S} = \begin{bmatrix} S(E, E) + 2(1-a) & S(E, \varphi E) & 0 \\ S(\varphi E, E) & S(\varphi E, \varphi E) + 2(1-a) & |\sigma| \\ 0 & |\sigma| & S(X_0, X_0) + 2(a^2 - 1) \end{bmatrix}.$$

By [3, p. 370], $S(X_0, X_0) = 2(1 - c^2/4)$. By [5, Lemma 7.1], the sectional curvatures $K(X_0, X)$ and $K(X_0, \varphi X)$ are related by

$$K(X_0, X) - K(X_0, \varphi X) = -(\nabla_{X_0}\tau)(X, X) = 2\psi(X, X)$$

for any unit vector $X \in B$. Thus,

$$S(E, E) = S(\varphi E, \varphi E) + 2\psi(E, E).$$

By polarization,

$$S(E, \varphi E) = \psi(E, \varphi E).$$

For,

$$S(E + \varphi E, E + \varphi E) - S(\varphi E - E, \varphi E - E) = 2\psi(E + \varphi E, E + \varphi E),$$

from which $4S(E, \varphi E) = 2[\psi(E, E) + \psi(E, \varphi E) + \psi(\varphi E, E) + \psi(\varphi E, \varphi E)] = 4\psi(E, \varphi E)$ since by [3, Proposition 1], $\text{trace } \psi = 0$ and φ is symmetric with respect to ψ .

Now, the scalar curvature $r = \text{trace } S$, so that

$$\begin{aligned} r &= S(E, E) + S(\varphi E, \varphi E) + S(X_0, X_0) \\ &= S(E, E) + S(E, E) - 2\psi(E, E) + 2\left(1 - \frac{c^2}{4}\right) \end{aligned}$$

from which

$$S(E, E) = \frac{r}{2} + \frac{c^2}{4} - 1 + \psi(E, E)$$

and

$$S(\varphi E, \varphi E) = \frac{r}{2} + \frac{c^2}{4} - 1 - \psi(E, E).$$

It follows that

$$\tilde{S} = \begin{bmatrix} \frac{r}{2} + \frac{c^2}{4} + 1 - 2a + \psi(E, E) & \psi(E, \varphi E) & 0 \\ \psi(\varphi E, E) & \frac{r}{2} + \frac{c^2}{4} + 1 - 2a - \psi(E, E) & |\sigma| \\ 0 & |\sigma| & 2\left(1 - \frac{c^2}{4}\right) + 2(a^2 - 1) \end{bmatrix}.$$

The inequalities $c < 2a$ and (4) ensure that $\tilde{S} > 0$ at $x \in M$. Indeed, since $c^2 = \psi(E, E)^2 + \psi(E, \varphi E)^2$, $\tilde{S}(E, E) > 0$, and

$$\begin{aligned} \tilde{S}(E, E)\tilde{S}(\varphi E, \varphi E) - \tilde{S}(E, \varphi E)^2 &= \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a\right)^2 - \psi(E, E)^2 - \psi(E, \varphi E)^2 \\ &= \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a\right)^2 - c^2 > 0. \end{aligned}$$

Moreover,

$$\begin{aligned} & -|\sigma| \left\{ |\sigma| \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a \right) + \psi(E, E) \right\} \\ & + \left\{ 2\left(1 - \frac{c^2}{4}\right) + 2\left(a^2 - \frac{c^2}{4}\right) \right\} \left\{ \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a \right)^2 - c^2 \right\} \\ & > -|\sigma|^2 \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a + c \right) + 2\left(a^2 - \frac{c^2}{4}\right) \left\{ \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a \right)^2 - c^2 \right\} \\ & = \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a + c \right) \left\{ 2\left(a^2 - \frac{c^2}{4}\right) \left(\frac{r}{2} + \frac{c^2}{4} + 1 - 2a - c \right) - |\sigma|^2 \right\} > 0. \end{aligned}$$

This completes the proof. The last part is a consequence of a theorem of Hamilton [4].

COROLLARY. *Let M be a compact oriented 3-manifold with K -contact metric structure $(\varphi, X_0, \omega, g)$. If $r > -2$, then M admits a contact metric of positive Ricci curvature.*

REMARK. The quantity $r/2 + c^2/4 + 1$ in (4) is equal to $r^*/2$, where r^* is the generalized Tanaka-Webster scalar curvature defined in [5, p. 21].

REFERENCES

- [1] S. S. CHERN AND R. S. HAMILTON, On Riemannian metrics adapted to three-dimensional contact manifolds, Lecture Notes in Math., Springer-Verlag, Berlin and New York, Vol. 1111, 1985.
- [2] S. I. GOLDBERG, Nonnegatively curved contact manifolds, Proc. Amer. Math. Soc. 96 (1986), 651–656.
- [3] S. I. GOLDBERG AND G. TOTH, Torsion and deformation of contact metric structures on 3-manifolds, Tohoku Math. J. 39 (1987), 365–372.
- [4] R. S. HAMILTON, Three-manifolds with positive Ricci curvature, J. Diff. Geom. 17 (1982), 255–306.
- [5] S. TANNO, Variational problems on contact Riemannian manifolds, preprint.

DEPARTMENT OF MATHEMATICS UNIVERSITY OF ILLINOIS 1409 WEST GREEN STREET URBANA, ILLINOIS 61801 U.S.A.	AND	DEPARTMENT OF MATHEMATICS RUTGERS UNIVERSITY CAMDEN, N.J. 08102 U.S.A.
---	-----	---

AND

DEPARTMENT OF MATHEMATICS
 AND STATISTICS
 QUEEN'S UNIVERSITY
 KINGSTON, ONTARIO K7L 3N6
 CANADA